Clinical Data Summary

This assessment synthesizes a transthoracic echocardiogram (TTE, October 21, 2025), two 12-lead ECGs (including one dated September 3-4, 2025), and a lipid panel for a 72-year-old male with a 40-year history of vigorous, near-daily exercise and 20 years of statin therapy.

Lipid Panel

Marker	Result	Normal Range	Interpretation
Total cholesterol	129 mg/dL	100-199	Excellent
Triglycerides	47 mg/dL	0-149	Excellent
HDL cholesterol	78 mg/dL	greater than 39	Excellent — very high, cardioprotective
VLDL cholesterol	11 mg/dL	5-40	Normal
LDL cholesterol, calculated	40 mg/dL	0-99	Excellent — very aggressive control

This lipid profile is exceptional by any standard, reflecting two decades of effective statin therapy. LDL of 40 mg/dL is well below even the most aggressive secondary-prevention targets (typically less than 70, or less than 55 for very-high-risk patients), and an HDL of 78 mg/dL is strongly cardioprotective. Long-term statin use in patients over 70 is well supported by clinical evidence: meta-analyses of trials including JUPITER and HOPE-3 show a 26 percent relative risk reduction in nonfatal MI, nonfatal stroke, or cardiovascular death in patients over 70, and population studies of continuous statin therapy in older adults show a 21 percent relative risk reduction in cardiovascular disease for ages 75-84. This lipid result essentially removes atherosclerotic/lipid-driven coronary risk as a major concern, though it does not eliminate risk from other structural or rhythm-based factors detailed below.[1][2][^3]

Electrocardiogram Findings

Both ECGs are concordant and show:

- Atrial flutter with second-degree AV block at a variable conduction ratio, ventricular rate 39-45 bpm (bradycardic)
- Left axis deviation, consistent with left anterior fascicular block (LAFB)
- Incomplete right bundle branch block (RBBB)
- A pattern read as "anteroseptal infarct" / "old anterior infarct," with slight inferior repolarization disturbance flagged as a consideration for ischemia
- Prolonged QT/QTc (492/396 ms)
- Overall interpretation: Abnormal ECG

Echocardiogram Findings

Parameter	Value	Reference	Interpretation
LV ejection fraction	59%	Normal (≥55%)	Normal systolic function
LV wall thickness (septum/posterior wall)	1.1 cm / 1.1 cm	Upper-normal to mild	Mild concentric LVH
LV mass (2D)	250 g	Elevated for BSA	Increased LV mass
LV chamber size (LVIDd)	5.7 cm	4.2-5.9	Normal, upper range
Right ventricle size	Moderately enlarged	—	Enlarged
Left atrium (diameter/volume index)	5 cm / LAVi 41.8 mL/m²	3-4 cm / 16-34 mL/m²	Moderately enlarged
Right atrium (area)	29.5 cm²	10-18 cm²	Severely enlarged
Aortic valve	Mild sclerosis, mild regurgitation	—	Mild, non-stenotic
Mitral/tricuspid/pulmonic valves	Trace-to-mild regurgitation	—	Non-significant
Pericardium	No effusion	—	Normal
Rhythm during study	Atrial flutter, HR 45-50 bpm	—	Confirmed arrhythmia

Integrated Cardiovascular Assessment

Structural Findings: Athletic Remodeling vs. Pathology

The echocardiographic picture — mild concentric LV hypertrophy, increased LV mass, and enlargement of both atria and the right ventricle, all with preserved biventricular systolic function — is a textbook description of exercise-induced cardiac remodeling ("athlete's heart"), a well-documented adaptation to decades of sustained endurance training. Four decades of near-daily vigorous exercise is precisely the exposure associated with this pattern.[4][5][^6]

Key supporting evidence: a meta-analysis of 46 studies (6,806 athletes) established an upper limit of normal right atrial area of 23 cm² in athletes, nearly double the general-population threshold, because sustained increases in cardiac output during training produce balanced, proportional enlargement of all four chambers. Studies of endurance athletes similarly show left atrial enlargement in up to 20 percent of long-term competitors, with more marked enlargement in about 2 percent, generally interpreted as benign when atrial function (not just size) remains normal. Right heart enlargement, including RV and RA, is often the most prominent feature in endurance athletes specifically, and can appear disproportionate to left heart changes without indicating pathology. Notably, the echo report describes left ventricular diastolic parameters as consistent with normal left atrial pressure, an important reassuring detail: the atrial enlargement is volumetric (from decades of stroke-volume loading), not pressure-driven (as would occur with hypertension, valvular disease, or heart failure).[7][8][5][6][9][4]

That said, the right atrium's severe enlargement (29.5 cm^2, well above even the athletic-population ceiling of 23 cm^2) is more pronounced than typical athletic remodeling and warrants attention rather than reassurance alone. Isolated or disproportionate right heart enlargement is a recognized area of diagnostic overlap between benign athletic adaptation and pathologic conditions (e.g., arrhythmogenic right ventricular cardiomyopathy, pulmonary hypertension, or atrial cardiomyopathy), and guidance specifically flags that isolated RV dilation not matched by proportional LV change should raise clinical suspicion. In this case, pulmonary artery pressures were not elevated (no evidence of pulmonary hypertension) and RV/LV enlargement appear roughly proportional, which is reassuring, but the magnitude of right atrial enlargement — plus the presence of atrial flutter itself — fits a related and increasingly recognized concept: exercise-associated atrial cardiomyopathy.[8][9]

Atrial Cardiomyopathy and the Athlete's Paradox

A growing body of literature identifies "atrial cardiomyopathy" (AtCM) as a downstream consequence of the same chronic volume overload that produces athletic atrial enlargement. Endurance athletes, despite having structurally "adapted" hearts, show a demonstrably higher prevalence of atrial fibrillation/flutter than age-matched non-athletes, a paradox attributed to atrial stretch, fibrosis, and altered autonomic tone from decades of training. This directly bears on your case: the presence of atrial flutter is not incidental to the atrial enlargement — it is mechanistically linked to it. The same 40 years of vigorous training that produced the favorable lipid and hemodynamic profile may also be a contributing driver of the atrial arrhythmia captured on both ECGs and during the echo study.[^10]

The Arrhythmia: Atrial Flutter with High-Grade AV Block

The most clinically significant and time-sensitive finding is atrial flutter with second-degree AV block and a resulting ventricular rate of 39-45 bpm — a high-grade AV block presentation that is distinct from ordinary atrial flutter (which typically produces tachycardia near 150 bpm). This combination changes clinical priorities substantially:[^11]

- In the absence of AV-nodal-blocking medications (beta-blockers, calcium channel blockers, digoxin, amiodarone), a high-grade block with flutter most often reflects intrinsic AV nodal or infranodal conduction disease, which becomes more common with age and can be a marker of underlying degenerative conduction system disease.[12][11]
- Second-degree AV block of the Mobitz II type, or any high-grade block producing symptomatic or sustained bradycardia below roughly 40 bpm, is generally considered a Class I indication for permanent pacemaker implantation, because these blocks are prone to progression to complete heart block.[13][14]
- Once a pacemaker is in place, rate control of the flutter with beta-blockers or calcium channel blockers becomes safe; before pacing, these drugs are contraindicated because they would worsen the bradycardia.[^14]
- Catheter ablation of the cavotricuspid isthmus is a definitive, often curative treatment for typical atrial flutter and has been used successfully in similar elderly presentations with slow ventricular response, sometimes resolving both the flutter and the bradycardia without a pacemaker.[^12]

This arrhythmia also carries thromboembolic (stroke) risk that must be formally assessed. Atrial flutter is treated identically to atrial fibrillation for stroke risk stratification using the CHA_2DS_2-VASc score. Based on the information available (age 72, male), a rough calculation gives a baseline of 1 point for age 65-74, which alone crosses the guideline threshold recommending consideration of anticoagulation in men (score of 1 or higher). If any additional risk factors are present (hypertension, diabetes, vascular disease, prior stroke/TIA, or heart failure — each adds 1-2 points), the score rises further, and at 2 or more points anticoagulation is a Class I recommendation with strong evidence. Annual stroke risk climbs from roughly 0.6-1.5 percent at a score of 1 to 2.2-2.9 percent at a score of 2, and continues rising thereafter. This is a critical, actionable item to confirm with your treating cardiologist/electrophysiologist if anticoagulation has not already been addressed.[15][16][17][18][^19]

The ECG Pattern Read as "Anteroseptal/Old Anterior Infarct"

The ECG's automated and physician-reviewed interpretation flags an anteroseptal or "old anterior" infarct pattern along with left anterior fascicular block and incomplete RBBB. This combination deserves careful contextualization rather than alarm: LAFB and RBBB both alter the QRS axis and can produce poor R-wave progression or Q-wave-like patterns in the anterior/septal leads that mimic infarction on automated and even expert ECG reads, independent of any actual myocardial scar. The echocardiogram is the more decisive test here, and it directly addresses this question — the echo explicitly reports normal segmental wall motion with no regional wall motion abnormality and preserved global systolic function (EF 59%), which is generally the most sensitive bedside indicator of prior significant infarction. A true anterior wall myocardial infarction of the sort implied by "old anterior infarct" almost always produces a regional wall motion abnormality (hypokinesis, akinesis, or aneurysm) in the corresponding LV segment; its absence here is reassuring evidence against a hemodynamically significant prior infarct, though it does not rule out a small, non-transmural, or remote silent event. This discrepancy between the ECG pattern and normal wall motion should be discussed directly with the cardiologist, as it may simply reflect conduction-related ECG changes (from LAFB/incomplete RBBB) rather than true infarction.[^9]

Prolonged QTc and AV Nodal Disease

A QTc of 396 ms is within normal limits (the 492 ms figure reflects the uncorrected QT at a very slow heart rate, which mathematically prolongs raw QT without necessarily indicating true repolarization abnormality). This is a byproduct of the bradycardia from the AV block rather than a standalone risk factor, but it reinforces that the conduction system — not the myocardium or coronary arteries — is the dominant abnormality in this case.

Synthesis: Overall Risk Profile

Domain	Finding	Risk Implication
Atherosclerotic/lipid risk	LDL 40, HDL 78, TG 47	Very low — 20 years of effective statin therapy
Coronary/ischemic risk	Normal wall motion, EF 59%, no regional WMA	Low — no echo evidence of significant infarct despite ECG pattern

Domain	Finding	Risk Implication
Structural remodeling	Mild concentric LVH, biatrial and RV enlargement	Largely consistent with athletic heart; RA enlargement more pronounced than typical
Valvular disease	Mild AV sclerosis/regurgitation, trace MV/TV/PV regurgitation	Low — no stenosis, no significant dysfunction
Arrhythmia/conduction	Atrial flutter with 2nd-degree AV block, bradycardia 39-45 bpm	High priority — needs EP evaluation, possible pacemaker/ablation
Thromboembolic/stroke risk	Atrial flutter present, age 72	Elevated — CHA_2DS_2-VASc likely ≥1, anticoagulation should be addressed

Taken as a whole, this is a cardiovascular profile with two very different faces. On one axis — atherosclerotic and myocardial disease — the outlook is excellent: two decades of statin-driven lipid control have produced a lipid panel better than most healthy young adults, and the echocardiogram shows preserved, normal pump function with no wall motion abnormality despite an ECG pattern suggestive of prior infarct. On the other axis — the cardiac conduction system and atrial substrate — there is a real and active abnormality: atrial flutter with high-grade AV block and pronounced right atrial enlargement, a combination that is plausibly linked to decades of endurance exercise but that nonetheless requires electrophysiological management, most likely pacemaker evaluation and/or catheter ablation, along with formal stroke-risk-based anticoagulation decision-making [c, cite:20]. Vigorous lifelong exercise appears to explain the favorable lipid/vascular findings only indirectly (through general cardiovascular fitness and metabolic health) while directly explaining the atrial and right-heart chamber enlargement; it is also a recognized, if incompletely understood, contributor to the atrial arrhythmia burden seen in long-term endurance athletes. None of these findings should be interpreted as reflecting poorly managed cardiovascular health — rather, they represent the specific, well-described tradeoff profile of an aging endurance athlete: outstanding vascular/lipid health paired with an atrial/conduction system that has been volume-loaded for four decades and is now manifesting rhythm disease that needs its own dedicated, likely procedural, management.[10][14][^12]

References

1. Study finds statins reduce the relative risk of cardiovascular ... - Researchers provided comprehensive evidence of the benefits of statin use in elderly patients, addre...
2. Primary Prevention With Statin Therapy in the Elderly | Circulation
3. Statin Therapy for Primary and Secondary Prevention in Older ... - Although statin therapy is well established to prevent atherosclerotic vascular disease (ASCVD) even...
4. Atrial Enlargement in the Athlete's Heart - American College of Cardiology
5. Athlete heart syndrome | Radiology Reference Article | Radiopaedia.org - Athlete heart syndrome refers to adaptations in both cardiac structure and function seen in people e...
6. Cardiac Imaging In Athletes - PMC - NIH - Athletic heart syndrome refers to the physiological and morphological changes that occur in a human ...

7. Atrial Enlargement in the Athlete's Heart: Assessment of ... - by F D'Ascenzi · 2018 · Cited by 113 — Atrial size is often increased in competitive athletes. • In ...
8. The Athlete's Right Heart
9. Athlete's Heart Section 4 - ... right heart often increases proportionally to left heart enlargement. In endurance athletes, rig...
10. Atrial cardiomyopathy in endurance athletes - by LW Spencer · 2024 · Cited by 14 — Changes in atrial size and function are frequently observed in ...
11. Unusual Presentation of Atrial Flutter With Slow Ventricular Response - Atrial flutter is usually associated with tachycardia with a ventricular rate of 150 beats per minut...
12. Atrial Flutter with Variable High-Degree AV Block - medtigo Journal
13. High-grade atrioventricular block - PMC - Atrioventricular (AV) block is an AV conduction disorder that can manifest in various settings, with...
14. What is the best treatment approach for an older adult patient with ... - For an older adult with atrial flutter and AV block, permanent pacemaker implantation is the definit...
15. CHA_2DS_2-VASc score — The Cardiovascular - Stroke risk in non-valvular atrial fibrillation, to guide oral anticoagulation.
16. Outcomes in Patients With Atrial Fibrillation or Atrial Flutter by CHA2DS2-VASc Score - This cohort study uses Taiwan National Health Insurance database data to evaluate the incidence of i...
17. Stroke in atrial fibrillation: Review of risk stratification and preventive ... - Atrial fibrillation (AF) is a leading preventable cause of ischemic stroke for which early detection...
18. VASc for stroke risk stratification in patients with atrial fibrillation ... - PMC
19. CHA2DS2–VASc score - Wikipedia